

COVID-19 in Italy: impact of containment measures and prevalence estimates of infection in the general population

Carlo Signorelli^{1,2}, Thea Scognamiglio³, Anna Odone^{1,4}

¹School of Medicine, Vita-Salute San Raffaele University, Milan; ²Past-President Italian Society of Hygiene, Preventive Medicine and Public Health (SIItI); ³Johns Hopkins Center for Humanitarian Health; ⁴IRCCS San Raffaele Scientific Institute, Milan

Abstract: Since the beginning of the COVID-19 epidemic in Italy, the Italian Government implemented several restrictive measures to contain the spread of the infection. Data shows that, among these measures, the lockdown implemented as of 9 March had a positive impact, in particular the central and southern regions of Italy, while other actions appeared to be less effective. When the true prevalence of a disease is unknown, it is possible estimate it, based on mortality data and the assumptive case-fatality rate of the disease. Given these assumptions, the estimated period-prevalence of COVID-19 in Italy varies from 0.35% in Sicily to 13.3% in Lombardy.

Key words: COVID-19, containment, prevalence, mortality

On April 7th, 2020, the number of notified COVID-19 cases in Italy is above 135,000, with almost 17,000 deaths (1). Italy, second only to the US for the number of COVID-19 notified cases, but first for of deaths, is facing an unedited challenge (2). The epidemic is disproportionately hitting some northern areas, pushing the Italian National Health Service capacity of some areas to their limits (3-5).

Lessons from China's successful battle against COVID-19 show how containing measures including cases isolation, contact tracing, and quarantine and mitigation measures, including general lockdown and social (or personal) distancing, seem to have worked (6-8).

As COVID-19 was first reported in Italy (23rd January 2020), the Government has progressively introduced restrictive measures (1,9). The most relevant actions taken by the Government from 25 January to 21 March 2020, are described in Table 1. Prevention measures, taken in late January, such as health-check-points in airports and a flight ban from China, rapidly escalated when the first autochthonous cases were detected in the area of Casalpusterlengo, Castelnuovo


d'Adda and Codogno in the Lombardy Region (21 February) (Figure 1).

Containment measures at first had been the self-isolation of infected patients and the creation of a "red-zone" in a limited area. Afterward, the Government strategy changed into a severe mitigation response, up to the lockdown to all the country. Other mass-measures were introduced, such as a strong recommendation to avoid at-risk behaviors and the suspension of all non-essential businesses in the entire nation.

The proportion of the infected population in Italy was estimated based on available published Italian data. This preliminary analysis did not consider infections notified within the national surveillance system to avoid biases due to the heterogenous proportion of population tested in different regions (1). The number of official deaths was considered to be less biased; therefore, we included an adjustment in our model due to the undiagnosed deaths, especially in the first phase of the epidemic (10). The average increase of 20% in the number of deaths was introduced accordingly to a recent Italian specific report (11). The adjusted case-

Table 1. Health protection measures against COVID-19 in Italy, 25 January- 23 March 2020

<i>Date</i>	<i>Public health measure implemented</i>	<i>Place</i>	<i>Authority</i>
25/01	Health checkpoints for passengers coming from China or from areas where one sustained autochthonous transmission of the new Coronavirus has occurred.	Airports, Italy	Ministry of Health
30/01	Air traffic from China is banned	Airports, Italy	Government
21/02	Mandatory supervised quarantine for 14 days for all individuals who have come into close contact with confirmed cases of disease; Mandatory communication to the Health Department from anyone who has entered Italy from high-risk of COVID-19 areas, followed by quarantine and active surveillance .	Public Health department in identified areas	Ministry of Health
23/02	Red zones: adoption of an adequate and proportionate containment and management measures in areas with >1 person positive to COVID-19 with unknown source of transmission.	11 municipalities in Lombardy Region	Government
23/02	Suspension of all public events or open to the public, of any nature; Schools (all levels), public places, gyms, and other places of aggregation	5 Regions in Northern Italy	Ministry of Health
02/03	Proposal to extend the “red zone” to three additional municipalities in the provinces of Bergamo and Brescia from local authorities	Three municipalities in Lombardy Region	(not adopted)
08/03	The “national” Red Zone: avoid any movement of people except for motivated by proven work needs or situations of necessity (health, food and assistance); public and private employers should encourage to use days of ordinary leave and holidays and smart working; closing of ski facilities; limit travel and activities and sanitization measures and reduce close contacts	Lombardy Region (and other affected areas in 5 additional regions)	Government
11/03	Suspension of all business activities; Suspension of all commercial activities non-indispensable for production. Maximum use by companies of smart-working methods for activities that can be performed at home or remotely. Sanitation of workplace areas.	Italy	Government
23/03	Extension of the ban on non-indispensable activities. A list of 80 authorized activities is published. The ban extends limitations on individual freedom and on other business activities that were not explicitly closed by the previous measures.	Italy	Government

**Figure 1.** First “Red Zone” Area (February 23rd)

fatality rate for Europe (0.85%) was derived considering the averaged estimates of three reports, two carried out in UK and the other in Italy (12-14). From these parameters, we obtained a model to estimate the current period prevalence in the general population of the 19 Italian regions and the 2 autonomous provinces (Table 2). These data vary among Italian regions from 0.35% infection proportion in Sicily and Basilicata to 11.2% in the Lombardy region and are a useful tool to inform the planning of further containment measures in different geographical areas.

Conclusions

The epidemic showed an evident spread. If we analyze these trends, we can outline the following, preliminary considerations:

a) The measures to suspend flights from China (which were not adopted by other EU countries) and air-

ports' checkpoints with *thermoscan* did not have a significant effect in containing the epidemic.

- b) The implementation of a “red zone” in Lombardy effectively contained the spread of the infection within that area. On the other hand, the “red zone” measure did not have the same effect outside that area. In fact, three of the neighboring provinces (Bergamo, Brescia, and Piacenza) recorded the highest incidence rates in the weeks following the establishment of the red zones (1,5). Perhaps these actions were adopted too late (or considering a too small area) when the virus had already spread for several days without notice of it.
- c) The failure to establish a second “red zone” near Bergamo in the Municipalities of Alzano and Nembro (Figure 2), despite the proposal of local authorities (on March 3rd), led to a dramatic outbreak with about 10,000 cases in Bergamo with over 1,000 death toll and similar figures in the

Table 2. Population, Number of deaths and estimates of infection period prevalence in the Italian Regions and Autonomous Provinces

<i>Regions and Autonomous Provinces</i>	<i>Population M (mil)</i>	<i>N. of deaths</i>	<i>Period Prevalence</i>
Piemonte (Piedmont)	4.4	1.319	4.3%
V.d'Aosta (Aosta Valley)	0.13	100	11.2%
Lombardia (Lombardy)	10.1	9.484	13.3%
Prov. Aut. Bolzano	0.52	174	4.7%
Prov. Aut. Trento	0.54	244	6.4%
Veneto	4.9	695	2.0%
Friuli-Venezia Giulia	1.3	164	1.8%
Liguria	1.6	620	5.6%
Emilia-Romagna	4.5	2.180	6.9%
Toscana (Tuscany)	3.7	369	1.4%
Umbria	0.88	49	0.78%
Marche	1.5	630	5.8%
Lazio	5.9	238	0.57%
Abruzzo	1.3	172	1.9%
Molise	0.31	13	0.60%
Campania	5.8	216	0.53%
Puglia (Apulia)	4.0	209	0.73%
Basilicata	0.56	14	0.35%
Calabria	1.9	60	0.43%
Sicilia (Sicily)	5.0	125	0.35%
Sardegna (Sardinia)	1.6	52	0.45%
Total Italy	60.4	17.127	4.0%



Figure 2. Missed “Red Zone” Area (March 3rd), from Regione Lombardia & Corriere della Sera




Figure 3. Trend of overall new notified cases in Italy (Protezione civile, 7,4,2020)¹

neighboring areas (Brescia and Piacenza) (5).

- d) General mitigation measures seem to be effective to flatten the epidemic curve of new notified infections (Figure 3) with more effect in controlling the epidemic in the central-southern than in the Northern regions (1).

The difference was in the time when the mass-measures were adopted. At the time of the national lock-down, the central-southern regions only had few circulating cases. Timing appeared to be a crucial factor in determining the effect of mitigation measures (Figure 4).

Overall, containment measures (red zones) and mitigation (general lockdown) can be effective if taken at an early stage of the epidemic and on large areas. Also, the community management of suspects, contacts, and cases could alleviate hospital burden and perhaps even improve disease prognosis.

The post-epidemic phase might benefit from the availability of forthcoming antibody serological tests. The benefit could be substantial for a large part of the country population, but especially central and southern Italy, that would not yet be infected, as shown by our estimates.



Figure 4. Estimate of Period Prevalence of infected people in the Italian regions as at 7 April, 2020

Funding: This paper is a preliminary activity among the EU Project n. 101003562 “Three Rapid Diagnostic tests (Point-of-Care) for COVID-19 Coronavirus, improving epidemic preparedness, and foster public health and socio-economic benefits - CORONADIX” (Task 7.1) supported by the European Commission (Horizon 2020, H2020-SC1-PHE-CORONAVIRUS-2020).

Conflict of interest: Each author declares that he or she has no commercial associations (e.g. consultancies, stock ownership, equity interest, patent/licensing arrangement etc.) that might pose a conflict of interest in connection with the submitted article

References

1. Dipartimento della Protezione Civile. COVID-19 Italia - Monitoraggio della situazione. Aggiornato al 7/4/2020. <http://www.protezionecivile.gov.it/> (accessed 08.04.20)
2. European Centre for Disease Prevention and Control (ECDC). Situation update 6 april 2020. <https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases> (accessed 08.04.20)
3. Lazzerini M, Putoto G. COVID-19 in Italy: momentous decisions and many uncertainties. *Lancet Glob Health*. 2020 Mar 18. pii: S2214-109X(20)30110-8
4. Signorelli C, Odone A, Oradini-Alacreu A, Pelissero G. Universal Health Coverage in Italy: lights and shades of the Italian National Health Service which celebrated its 40th anniversary. *Health Policy*. 2020 Jan;124(1):69-74.
5. Boccia S, Ricciardi W, Ioannidis JPA. What Other Countries Can Learn From Italy During the COVID-19 Pandemic. *JAMA Intern Med*. 2020 Apr 7. doi: 10.1001/jamainternmed.2020.1447. [Epub ahead of print]
6. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 Epidemic?. *The Lancet*. Published: 9 March 2020.
7. WHO. Pandemic Influenza Preparedness and Response: A WHO Guidance Document. Geneva: World Health Organization; 2009. 4, The WHO pandemic phases. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK143061/>
8. Lau H, Khosrawipour V, Kocbach P, Mikolajczyk A, Schubert J, Bania J, Khosrawipour T. The positive impact of lockdown in Wuhan on containing the COVID-19 outbreak in China. *J Travel Med*. 2020 Mar 17. pii: taaa037. doi: 10.1093/jtm/taaa037.
9. Gazzetta Ufficiale della Repubblica Italiana, Raccolta degli atti recanti misure urgenti in materia di contenimento e gestione dell'emergenza epidemiologica da COVID-19
10. ISTAT L'andamento dei decessi del 2020. Dati anticipatori sulla base del sistema ANPR. Nota esplicativa 2020 Mar 31
11. UCSC. Coronavirus, le stime dei ricercatori: "In Italia almeno 2800 morti non dichiarati" (to be completed)
12. Oke J, Heneghan C. Global Covid-19 Case Fatality Rates. The Centre for Evidence-Based Medicine develops, promotes and disseminates better evidence for healthcare (CEBM). <https://www.cebm.net/covid-19/global-covid-19-case-fatality-rates/> (accessed 07.04.20)
13. Ferguson N, et al. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand. 16 Mar 2020; doi.org/10.25561/77482
14. Villa M. La letalità italiana tra apparenza e realtà. ISPI, 27 3 2020. <https://www.ispionline.it/it/pubblicazione/coronavirus-la-letalita-italia-tra-apparenza-e-realta-25563>

Received: 7 April 2020

Accepted: 8 April 2020

Correspondence:

Carlo Signorelli

School of Public Health (Igiene e medicina preventiva)

University Vita-Salute San Raffaele

Via Olgettina, 58

20132 - Milano (Italy)

E-mail: signorelli.carlo@hsr.it